Meta-analysis

Impact of resection vs respect techniques on left ventricular function after mitral valve repair: a systematic review and meta-analysis

Casey Christiany¹, Victor Jesron Nababan¹, Darma Satria²

¹Department of Thoracic, Cardiac, and Vascular Surgery, BP Batam Hospital, Batam, Riau Island, Indonesia

Abstract

Objective: Mitral valve disease, ranging from degenerative to infective origin, is one of the most prevalent left heart diseases globally and affects a large number of individuals. Conventionally, surgical repair has grown to become the treatment of choice, with two main techniques being resection and respect. Current literature has yet to address which technique is superior to the other, particularly in regard to left ventricular (LV) function.

Methods: We performed a systematic review and meta-analysis on three databases with a primary outcome of LV function alongside its' parameters, and a secondary outcome of repair durability, mitral valve gradient, and mortality rates. Meta-analysis was performed using random effects, and results were displayed in forest plots. Risk of bias was conducted using the Newcastle-Ottawa Scale.

Results: Six retrospective studies were included, evaluating a total of 3376 patients. Pooled results showed that LV function were preserved equally in both groups, showing no statistically significant differences. The respect group had slightly lower mortality rates in comparison to the resect group, and the repair success rate showed a slight superiority in the respect group. Significant heterogeneity was observed on left atrial diameter (LAD) measurement, indicating variability. The overall differences in LV function coming from both techniques appear intangible. Plenty of consideration must be made beyond LV function in determining which repair technique should be performed on a patient.

Conclusion: This study demonstrated that both the resect and respect techniques were found to be equally excellent in preserving left ventricular function after mitral valve repair.

Key words: Mitral valve repair, left ventricular function, respect, surgical repair

(Heart Vessels Transplant 2025: 9: doi:10.24969/hvt.2025.605)

Highlights

This systematic review and meta-analysis compared the resect and respect techniques in mitral valve repair. Both approaches demonstrated equivalent efficacy in preserving left ventricular function, with no significant differences in LVEF, LVEDD, or LVESD across 3,376 patients from six retrospective studies. Secondary outcomes such as repair success and mortality showed inconsistent findings, underscoring that neither method was clearly superior. These results suggest that surgical strategy should be individualized based on valve anatomy, surgeon expertise, and patient-specific considerations, while future prospective trials are warranted to clarify long-term outcomes.

Address for Correspondence: Casey Christiany, Department of Thoracic, Cardiac, and Vascular Surgery, BP Batam Hospital, Batam, Riau Island, Indonesia

E-mail: caseychristiany03@gmail.com Phone: +6287884965742

ORCID: Casey Christiany - 0009-0001-1559-2147, Victor Jesron Nababan - 0009-0007-6973-6672,

Darma Satria - 0009-0008-0133-9184

Citation: Christiany C, Jesron Nababan VJ, Darma S. Impact of resection vs respect techniques on left ventricular function after mitral valve repair: a systematic review and meta-analysis. Heart Vessels Transplant 2025: 9: doi:10.24969/hyt.2025.605

Received: 29.09.2025 Revised: 31.10.2025 Accepted: 31.10.2025 Copyright ©2025 Heart, Vessels and Transplantation

²¹Department of Thoracic, Cardiac, and Vascular Surgery, Syiah Kuala University, Banda Aceh, Aceh, Indonesia

Graphical abstract




Impact of resection vs respect techniques on left ventricular function after mitral valve repair: a systematic review and meta-analysis

Casey Christiany, Victor Jesron Nababan, Darma Satria, Batam, Riau Island, Banda Aceh, Aceh, Indonesia


Background

Mitral valve repair is the treatment of choice for degenerative mitral regurgitation



Results

Equivalent LV function between resection and respect techniques



Slightly lower mortality and better repair success with respect techniques

Considerable heterogeneity in LAD

Resection strategies



Respect strategies



Methods

Systematic Review and Meta-Analysis

- · Databases searched
- · Primary outcomes soang
- · Secondary outcomes
- · Regression model regreeemie
- · Risk of bias assessment

Conclusions

Both techniques are equally effective

Technique should be individualized



Introduction

Mitral valve disease remains as one of the most prevalent left heart valve diseases globally, with an incidence rising significantly as of recent (1). Developed nations, in particular, have experienced an increase in patients with degenerative mitral valve disease, and developing countries are still in the middle of battling against rheumatic heart diseases (2, 3). Among these diseases, mitral regurgitation (MR) is the most prominent pathology requiring surgical intervention, with recent studies estimating the prevalence of moderate-to-severe MR at 0.67%, and mitral valve prolapse (MVP) at 2.6% globally. These numbers, albeit small in percentages, make up a large number of individuals affected by significant mitral valve diseases (1, 4).

Conventionally, surgical repair has grown to become the treatment of choice for most patients with degenerative MR, with evidence proving better survival rates and long-term outcomes in comparison to mitral valve replacement (MVR) (5). In comparison to MVR, mitral valve repair has shown less operative risks; more ventricular function preserved, as well as reduced complications related to prosthetic valves. Current guidelines recommend surgical repair over replacement when feasible (5, 6).

Currently, two primary techniques are offered for mitral valve repair, being resection (or "resect") technique, and the respect technique. The resect approach focuses on the excision of the diseased leaflet tissue, followed by the reconstruction of the remaining viable tissue (7, 8). On the other hand, the respect approach utilizes artificial neochordaes to re-suspend prolapsed leaflet segments, which is aimed to preserve the leaflet anatomy and structure. Ultimately, both techniques aim to restore valve competence and durability, however, their repercussions on valve function and clinical outcomes, such as left ventricular (LV) performance, remains unknown (9). Studies have found contrasting answers, with one study stating how the respect approach is a better modality in comparison to the former in terms of permanent pacemaker implantation rates and mean gradients (10), and other studies asserting how the resect approach is more reproducible and therefore are more commonly used (11). Other trials have also remained inconclusive (12, 13).

Current literature has yet to address the consequences or improvements of these techniques to the LV function, durability, and patient outcomes.

This systematic review and meta-analysis aim to bridge literature gaps and directly compare the resection versus respect techniques in mitral valve repair, focusing on their impact on left ventricular function, as well as other clinical outcomes.

Methods

This study is a systematic review and meta-analysis performed based on the Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) 2020 guidelines (14). Our population included adult patients, categorized as individuals aged 18 and above undergoing mitral valve repair, with or without degenerative mitral valve disease, including stenosis and regurgitation. The intervention we chose to proceed with was the resecting technique, in comparison to the respecting technique. This study has two outcomes, with the primary outcome being changes in LV function, including left ventricular ejection fraction (LVEF), left ventricular end systolic diameter (LVESD), and left ventricular end diastolic diameter (LVEDD) after surgery. Our secondary outcome includes durability of repair, mitral valve gradient, and mortality rates after repair.

Literature searching was systematically performed in three databases up until July 29, 2025, including PubMed, Science Direct, and Google Scholar, with keywords tailored to each site as seen in Table 1. The general search terms included "Mitral Valve Repair"," Resecting technique OR Resection technique", and "Left Ventricular Function OR LVEF OR LVESD OR LVEDD". We made use of available Boolean operators when necessary to narrow down or broaden results. Searching and screening were performed by two independent reviewers, with a third reviewer to mediate any discrepancies when needed.

Our inclusion criteria included articles published from the past 10 years (2015-2025), fit into our population, intervention, comparison, and outcomes, reporting at least one LV function parameter pre- and post-surgery, with a follow up duration of at least 3 months. Our exclusion criteria included case reports, MVRs, studies without a comparison group, and studies whose full texts are irretrievable. The selected studies are then summarized into one table, with headings including but not limited to: (1) author; year of publication, (2) study design, (3) country of origin, (3) number of subjects included, (3) age

(years, mean standard deviation), (4) patients' characteristics, (5) mean follow-up duration, and (6) study findings.

Table 1. Searching Strategy							
Database	Keywords	Hits					
PubMed	("mitral valve repair"[MeSH Terms] OR "mitral valve repair") AND ("resection technique" OR "resecting technique" OR "leaflet resection") AND ("respect technique" OR "respecting technique" OR "chordal replacement") AND ("left ventricular function" OR "LVEF" OR "LVESD" OR "LVEDD")						
Science Direct	("mitral valve repair") AND ("resecting technique" OR "leaflet resection") AND ("respecting technique" OR "chordal replacement") AND ("left ventricular function" OR "LVEF" OR "LVESD" OR "LVEDD")	37					
Google Scholar	("mitral valve repair") AND ("resecting technique" OR "leaflet resection") AND ("respecting technique" OR "chordal replacement") AND ("left ventricular function" OR "LVEF" OR "LVESD" OR "LVEDD")	243					

The article we selected discusses the resect and respect techniques. The resect technique is based on the principle of removing (resecting) the leaflet that has prolapsed or become redundant in order to restore the valve to its normal shape. Examples include quadrangular resection, triangular resection, and leaflet resection with sliding plasty. In the respect technique, the focus is on preserving all valve tissue without resection, repairing the prolapse using neochordae.

Pre- and post- data collected were then observed for both respect and resect groups respectively.

Statistical analysis

To determine effect sizes, a random effects model was performed, with results reported as pooled mean differences with a 95% confidence interval (CI). Heterogeneity was quantified through the I2 statistics, with a value of I2 of over 50% defined as considerable heterogeneity (15). Forest plots are displayed to visualize findings. All analyses were performed using Review Manager (RevMan 5.4). A p-value of 0.05 and below determines statistical significance.

A comprehensive risk of bias would also be performed on the studies using the Newcastle-Ottawa Scale (16), scrutinizing domains including the selection, comparability, and outcome of the studies. Similarly, two independent reviewers proceeded with the critical appraisal, and any disparities will be settled by a third reviewer.

Results

The PRISMA Flowchart for this study is shown in Figure 1. Our initial search yielded 289 records, with 18 duplicates removed before further screening. Overall, 184 records were then excluded further. We attempted to retrieve 87 studies; however, 76 studies were unable to be retrieved; hence, 11 studies were assessed for their eligibility. We further excluded 5 studies for the unavailability of their full texts and included 6 studies in this review (17–22).

Risk of bias assessment was performed using Newcastle-Ottawa Scale for Cohort studies with three studies showing overall very good risk of bias and three studies showing good risk of bias (Supplementary Table 1.). Three studies showed a participant selection bias.

Study characteristics

Tables 1 and 2 summarize the study characteristics and the main findings of the study. Six retrospective studies were included in this review, evaluating a total of 3376 patients who has undergone mitral valve repair with either resection or respect techniques. The studies were performed in either European countries (i.e., The Netherlands, Germany) or Asian countries (Singapore, Japan, China). The sample size of each study ranged from 72 to 2134 subjects, with mean ages ranging from 55.3 to 64 years old.




Figure 1. PRISMA Flowchart

Patients' characteristics varied from having degenerative MR, to having an isolated posterior mitral leaflet (PML). The follow up duration varied from as short as one month, to as long as 87.6 months (17–22). Generally, the LVEF was preserved in both groups over all of the included studies. The LVEF values ranged from 58 to 68 percent in both groups. Similar results were also seen in the left atrial diameter (LAD), LVEDD, and LVESD measurements. Ma et al. (18) showed nearly identical LAD values (44.37 mm in the Respect group, in comparison to 44.63 mm for Resection groups). This was also seen in the study performed by Wijngaarden et al., measuring 37mm for both groups (19).

Secondary outcomes are also shown in Table 2. Ma et al. (18) reported a low in-hospital mortality for both groups, with 1 patient (1.4%) for the respect group, and 0 patients (0%), whereas Cetinkaya et al. (17) showed a 0.4% mortality rate for the respect group and 1.9% mortality rate for the resect group. Imasaka et al. (20), on the other hand, observed no deaths throughout the study period. In terms of repair success rate, a 95.8% success rate in the respect group was observed by Cetinkaya et al. (17) in comparison to a 86.7% repair success rate for the resect group. Wijngaarden et al. (19) measured a mitral valve mean gradient, with the respect group having a mean gradient of 2.8 mmHg, and the resect group having a mean gradient of 2.9 mmHg.

Table 1. Summ Author; year	Study	Country	No. of	Age	Patients'	Mean	LVEF,	%
of	design	,	samples	(years)	characteristics	follow-up	Mean (SD)	
publication			(Respect/ Resect)	(, =,		(months)	Respect	Resect
Wijngaarden et al., 2022 ¹⁹	Retrosp. cohort	Netherlands	125 (43/82)	63 (12)	Isolated PML, robotic repair	42.0 (14.5)	58 (7)	57 (9)
Ma et al., 2021 ¹⁸	Retrosp. (matched)	China	317 (243/74)	Respect: 55.3 (11) Resect: 55.8 (12)	Degenerative MR, more multi-segment in Respect	32 (median)	67.86 (6.07)	67.79 (5.51)
Pfannmueller et al., 2021 ²¹	Retrosp.	Germany	2134 (1751/383)	Respect: 58.6 (12.4) Resect: 58.9 (13)	Minimally invasive MV repair for MR	73.2 (51.6)	63.3 (8.6)	63.8 (11.1)
Cetinkaya et al., 2019 ¹⁷	Retrosp. (propensity -matched)	Germany	526 (263/263)	Respect: 61.8 (12.1) Resect: 62.1 (12.1)	Isolated PML prolapse	57.6 (respect) / 87.6 (resect)	60.8 (7)	61.5 (7.8)
Chua et al., 2016 ²²	Retrosp.	Singapore	202 (98/104)	Respect: 56.9 (10.9) Resect: 57.1 (12.1)	Degenerative MR, isolated PML	73.2 (48)	63.7 (7.9)	61.9 (7.9)
Imasaka et al., 2015 ²⁰	Retrosp.	Japan	72 (30/42)	Respect: 62 (15.1) Resect: 64 (11.3)	Isolated PML prolapse	1	62.9 (8.9)	65.5 (9.9)

LVEF – left ventricular rejection fraction, MR – mitral regurgitation, MV – mitral valve repair, PML – posterior mitral leaflet, rertrosp. – retrospective, SD –standard deviation

Visualized meta-analysis findings of this study in forest plots (Fig. 2). Overall, the pooled results of the meta-analysis showed the absence of statistically significant (overall effects: p>0.05) differences in left atrial or ventricular measurements (LAD, LVESD, LVEDD) and ejection fraction (LVEF) between the two groups. A significant heterogeneity is seen in the LAD measurement (I2= 80%), which indicates variability over the included studies.

Discussion

Findings of this systematic review and meta-analysis have shown that no differences were seen between the two techniques in preserving LV function. This is universally seen through the LAD size, LVEDD and LVESD measurements, as well as LVEF. In terms of the secondary outcomes, repair success rate was observed by one study (17), which showed a slight superiority of the respect group in comparison to the resect group.

Table 2. Summ	ary of stud	y results							
Author; year	LAD, mm		LVEDD, m	m	LVESD,	mm	LVEF, %		Other outcomes
of	(Mean (SI	O))	(Mean (SI	O))	(Mean (SI	O))	(Mean (SI	O))	
publication	Respect	Resect	Respect	Resect	Respect	Resect	Respect	Resect	
Wijngaarden	37 (6)	37 (8)	51 (6)	49 (8)	33 (7)	32 (7)	58 (7)	57 (9)	MV mean gradient
et al., 2022 ¹⁹									<u>(mmHg):</u>
									Respect: 2.8 (1.5)
									Resect: 2.9 (1.1)
Ma et al.,	44.37	44.63	55.87	55.99	33.35	34.47	67.86	67.79	In-hospital
2021 ¹⁸	(6.93)	(7.03)	(6.10)	(6.66)	(4.84)	(5.52)	(6.07)	(5.51)	mortality:
									Respect: 1 (1.4%)
									Resect: 0 (0%)
Pfannmueller	NR	NR	NR	NR	NR	NR	63.3	63.8	NR
et al., 2021 ²¹							(8.6)	(11.1)	
Cetinkaya et	54.8	52.4	55.0	55.7	34.3	35.0	60.8 (7)	61.5	Repair success rate:
al., 2019 ¹⁷	(10.0)	(10.2)	(8.2)	(6.7)	(6.2)	(6.2)		(7.8)	Respect: 252/263
									(95.8%)
									Resect: 228/263
									(86.7%)
									Mortality:
									Respect: 1/263
									(0.4%)
									Resect: 5/263
									(1.9%)
Chua et al.,	48.8	52.5	56.0	57.5	33.4	34.2	63.7	61.9	NR
2016 ²²	(9.1)	(9.6)	(7.2)	(7.2)	(6.0)	(5.5)	(7.9)	(7.9)	
Imasaka et	NR	NR	57.8	55.4	35.5	32.7	62.9	65.5	No deaths observed
al., 2015 ²⁰			(6.6)	(7.2)	(6.4)	(7.0)	(8.9)	(9.9)	and the land of the land


LAD – left atrial dimension, LVEDD - left ventricular end-diastolic dimension, LVESD – left ventricular end-systolic dimension, MV – mitral valve, NR – not reported, SD – standard deviation

However, this was the opposite for the mortality rate parameter, where the resect group had a slightly higher mortality rate. These parameters were not pooled; hence, we are unable to draw a conclusion for the secondary outcomes.

These findings demonstrate that both techniques were similarly capable of producing excellent LV outcomes, which is well supported by recent meta-analyses, illustrating that both methods lead to similar rates of long-term mortality, recurrence, and need of repeat surgeries, after adjusting for patients' individual risk factors (13). Additionally, previous matched cohort studies have suggested that both approaches are associated with high survival rates, durable repair, as well as reverse remodeling of the left-ventricle. The same study, however, mentioned that a smaller LVEDD was seen among the respect group in comparison to the

resect group (23). A different study has also seen a higher postoperative LVEF among patients who had undergone the respect technique (24). Regardless, the overall differences in functional recovery coming from both techniques appear intangible; hence, ultimately, the safety and efficacy of either approach are equally robust.


In order to effectively select the most appropriate repair technique, plenty of consideration must be made beyond LV function alone. This includes anatomical architecture, surgeon's capabilities, and valve pathology tailored to each patient. Given the indistinguishable long-term results, the decision to respect or resect should, overall, aim to achieve leaflet mobility with highest durability, in addition to patients' personal preferences (10, 23).




Comparison of LAD Size

	Re	Respect			Resect			Mean Difference	Mean Difference		
Study or Subgroup	Mean [mm]	SD [mm]	Total	Mean [mm]	SD [mm]	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI		
Cetinkaya 2019	55	8.2	263	55.7	6.7	263	29.4%	-0.70 [-1.98, 0.58]			
Chua 2016	56	7.2	98	57.5	7.2	104	20.3%	-1.50 [-3.49, 0.49]			
Imasaka 2015	57.8	6.6	30	55.4	7.2	42	11.0%	2.40 [-0.81, 5.61]	+		
Ma 2021	55.87	6.19	243	55.9	6.66	74	23.6%	-0.03 [-1.74, 1.68]	-		
Wijngaarden 2022	51	6	43	49	8	82	15.6%	2.00 [-0.49, 4.49]	+-		
Total (95% CI)			677			565	100.0%	0.06 [-1.18, 1.29]	•		
Heterogeneity: Tau ² : Test for overall effect			4 (P =	(0.10); f² = 49	9%			_	-10 -5 0 5 10 Favours [Respect] Favours [Resect]		

Comparison of LVEDD Size



Comparison of LVESD Size



Comparison of LVEF Size

Figure 2. Forest plots on Resection vs Respect techniques on left ventricular function

Study limitations and directions for future research

This study has several strengths, including the inclusion of the most recent studies, as well as a specific focus on the objective parameters of LV function. However, this study does come with its' limitations, including study heterogeneity, which may lead to inconsistent results. Moreover, the studies included have slight differences in defining 'resect' and 'respect', which may also contribute to additional heterogeneity and variety among study methods. Furthermore, follow-up duration between studies were widely incongruous. A head-to-head randomized controlled trials with a longer follow-

up duration may help establish a more uniformed study with less confounding and heterogeneity; hence, establishing a more credible and applicable finding. Additionally, comparing the measurement of longitudinal LV function, such as strain imaging, may also strengthen current findings.

Conclusion

This study demonstrated that both the resect and respect techniques were found to be equally excellent in preserving left ventricular function after mitral valve repair.

As this study primarily included retrospective studies, more prospective studies should be conducted to enhance the reliability of these findings.

Peer-review: External and internal **Conflict of interest:** None to declare

Authorship: C.C., V.J.N., and D.S. equally contributed to the study and manuscript preparation, thus fulfilled authorship criteria

Acknowledgement and funding: None to declare Statement on A.I.-assisted technologies use: The author s did not use Al-assisted technologies in preparation of this manuscript

Data and material availability: Does not apply

References

- 1. Figlioli G, Sticchi A, Christodoulou MN, Hadjidemetriou A, Amorim Moreira Alves G, De Carlo M, et al. Global prevalence of mitral regurgitation: A systematic review and meta-analysis of population-based studies. J Clin Med 2025; 14: 2749.
- 2. Chen QF, Shi S, Wang YF, Shi J, Liu C, Xu T, et al. Global, regional, and national burden of valvular heart disease, 1990 to 2021. J Am Heart Assoc 2024; 13: e037991.
- 3. Simpson MT, Kachel M, Neely RC, Erwin WC, Yasin A, Patel A, et al. Rheumatic heart disease in the developing world. Struct Heart 2023; 7: 100219.
- 4. Melamed T, Badiani S, Harlow S, Laskar N, Treibel TA, Aung N, et al. Prevalence, progression, and clinical outcomes of mitral valve prolapse: a systematic review and meta-analysis. Eur Heart J Qual Care Clin Outcomes 2025; qcaf016.
- 5. Lazam S, Vanoverschelde JL, Tribouilloy C, Grigioni F, Suri RM, Avierinos JF, et al. Twenty-year outcome after mitral repair versus replacement for severe degenerative mitral regurgitation. Circulation 2017; 135: 410–22.
- 6. Makarious Laham M, Easo J, Szczechowicz M, Roosta-Azad M, Weymann A, Ruhparwar A, et al. Five-year follow-up of mitral valve repair versus replacement: a propensity score analysis. J Cardiothor Surg 2023; 18: 27.
- 7. Wisniewski AM, Sutherland GN, Strobel RJ, Young A, Norman AV, Quader M, et al. Mitral valve repair in a regional quality collaborative: Respect or resect? JTCVS Techniques 2024; 24: 66–75.

- 8. Perier P, Hohenberger W, Lakew F, Diegeler A. Prolapse of the posterior leaflet: resect or respect. Ann Cardiothorac Surg 2015; 4: 273–7.
- 9. Calafiore A, Castellano G, Guarracini S, Di Marco M, Totaro A, Sacra C, et al. Respect or resect: a single strategy does not fit all. [cited 2025 Jul 29];Available from:

https://www.authorea.com/users/313145/articles/501766-respect-or-resect-a-single-strategy-does-not-fit-all 10. Sá MP, Cavalcanti LRP, Van den Eynde J, Amabile A, Escorel Neto AC, Perazzo AM, et al. Respect versus resect approaches for mitral valve repair: A study-level meta-analysis. Trends Cardiovasc Med 2023;33: 225–39. 11. Passos L, Aymard T, Biaggi P, Morjan M, Emmert MY, Gruenenfelder J, et al. Midterm outcomes of minimally invasive mitral valve surgery in a heterogeneous valve pathology cohort: respect or resect? J Thorac Dis [homepage on the Internet] 2023 [cited 2025 Jul 29];15(6). Available from: URL: https://jtd.amegroups.org/article/view/76143

- 12. Chan V, Mazer CD, Ali FM, Quan A, Ruel M, de Varennes BE, et al. Randomized, controlled trial comparing mitral valve repair with leaflet resection versus leaflet preservation on functional mitral stenosis. Circulation 2020; 142: 1342–50.
- 13. Caldonazo T, Sá MP, Jacquemyn X, Van den Eynde J, Kirov H, Harik L, et al. Respect versus resect approaches for mitral valve repair: A meta-analysis of reconstructed time-to-event data. Am J Cardiol 2024; 213: 5–11.
- 14. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ 2021; n71.
- 15. Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. BMJ 2003; 327: 557–60.
- 16. Wells G, B Shea, D O'Connell, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Ottawa Hospital Research Institute 2000;
- 17. Cetinkaya A, Bär S, Hein S, Bramlage K, Bramlage P, Schönburg M, et al. Mitral valve repair for posterior leaflet prolapse: Long-term comparison of loop implantation vs resection. J Card Surg 2020; 35: 11–20.
- 18. Ma J, Liu J, Wei P, Yao X, Zhang Y, Fang L, et al. Quadrangular resection versus chordal replacement for degenerative posterior mitral leaflet prolapse. Ann Transl Med 2021;9(1):60–60.

- 19. van Wijngaarden AL, Tomšič A, Mertens BJA, Fortuni F, Delgado V, Bax JJ, et al. Mitral valve repair for isolated posterior mitral valve leaflet prolapse: The effect of respect and resect techniques on left ventricular function. J Thorac Cardiovasc Surg 2022; 164: 1488-97.e3.
- 20. Imasaka K, Tayama E, Tomita Y. Left ventricular performance early after repair for posterior mitral leaflet prolapse: Chordal replacement versus leaflet resection. J Thorac Cardiovasc Surg 2015; 150: 538–45.
- 21. Pfannmueller B, Misfeld M, Verevkin A, Garbade J, Holzhey DM, Davierwala P, et al. Loop neochord versus leaflet resection techniques for minimally invasive mitral valve repair: long-term results. Eur J Cardiothorac Surg 2021; 59: 180–6.

- 22. Chua YL, Pang PYK, Yap YP, Abdul Salam ZH, Chen YT. Chordal reconstruction versus leaflet resection for repair of degenerative posterior mitral leaflet prolapse. ATCS 2016; 22: 90–7.
- 23. Del Forno B, Tavana K, Ruffo C, Carino D, Lapenna E, Ascione G, et al. Neochordae implantation versus leaflet resection in mitral valve posterior leaflet prolapse and dilated left ventricle: a propensity score matching comparison with long-term follow-up. Eur J Cardiothorac Surg 2023; 64: ezad274.
- 24. Mazine A, Friedrich JO, Nedadur R, Verma S, Ouzounian M, Jüni P, et al. Systematic review and metaanalysis of chordal replacement versus leaflet resection for posterior mitral leaflet prolapse. J Thorac Cardiovasc Surg 2018; 155: 120-8.e10.

Study		Sele	ction		Compar a- bility		Total quality score	AHRQ stan- dard		
	Representativeness of the EC	Selection on the non-EC	Ascer- tainment of the E	Demonst- ration that OI was not present at start of study	Comparability of subjects	Assess- ment of outcome	Adequate FU duration	Adequacy of cohort FU		
Wijngaarden et al., 2022 ¹⁹	a(*)	a(*)	a(*)	b(*)	a(**)	a(*)	b(*)	a(*)	9	Very good
Ma et al., 2021 ¹⁸	a(*)	С	a(*)	b(*)	a(**)	a(*)	b(*)	a(*)	8	Good
Pfannmueller et al., 2021 ²¹	a(*)	С	a(*)	b(*)	a(**)	a(*)	b(*)	a(*)	8	Good
Cetinkaya et al., 2019 ¹⁷	a(*)	a(*)	a(*)	b(*)	a(**)	a(*)	b(*)	a(*)	9	Very good
Chua et al., 2016 ²²	a(*)	С	a(*)	b(*)	a(**)	a(*)	b(*)	a(*)	8	Good
lmasaka et al., 2015 ²⁰	a(*)	a(*)	a(*)	b(*)	a(**)	a(*)	b(*)	a(*)	9	Very good

E- exposure, EC – exposed cohort, FU-follow-up, OI – outcome of interest