

Risk factors for the development of congenital heart defects in Kyrgyz Republic

Sofia M. Shakhnabieva, Shakhshianam Sh. Khasanova, Tatiana I. Sologubova, Elena I. Kondrateva, Ainagul D. Niiazalieva, Yuri M. Kurmanbakeev

Medical Faculty, Kyrgyz-Russian Slavic University, Bishkek, Kyrgyzstan

Abstract

Objective: Congenital heart diseases (CHD) are structural abnormalities of the heart and/or great vessels that are present at birth. CHD is still the most common inborn defect with an approximate prevalence of 5–11 per 1000 live births and incidence of 1%. CHDs are caused by many factors, often a mix of genetic predispositions and environmental influences, though the specific cause is unknown for most cases.

The aim was to study the risk factors for congenital heart defects, to identify risk groups in whom CHD may occur.

Methods: We conducted a survey of parents of children with CHD (the main group, n=392) and parents of healthy children (the control group, n=100), followed by an analysis of the clinical, anamnestic, and medical-social characteristics of parents of children in both groups. The following groups of risk factors were identified: demographic factors, socio-hygienic factors, biological factors, and chemical factors.

Results: The most important risk factors for the development of CHD in children were bad habits of parents: maternal smoking and alcoholism (4.1 (0.2) vs. 1.0 (0.2), $p<0.001$, main group vs control; 1.5 (0.1) vs. 0.0, $p<0.001$, respectively), paternal smoking (57.1 (1.6) vs. 39.0 (5.9), $p<0.01$), paternal alcoholism (28.6 (0.3) vs. 2.0 (0.5) $p<0.001$); infectious and inflammatory diseases of the mother during pregnancy: acute respiratory viral infections (40.8 (1.5) vs. 12.0 (2.6); $p<0.001$, main vs. control group), intrauterine infections (20.1 (1.0) vs. 1.0 (0.2), $p<0.001$); as well as the associated use of antibacterial drugs. Conditions directly related to pregnancy (severe toxicosis, maternal age, gestosis of pregnant women, anemia) made a less noticeable but significant contribution to determining the risk. The increased risk of having a child with CHD was determined by a number of socio-economic factors: marital status, low-income families (28.8 (1.3) vs 7.0 (1.6) main vs control group, $p<0.001$) and low educational level of the mothers of patients (9.4 (0.5) vs. 7.0 (1.6), $p<0.001$).

Conclusion: Thus, the most important risk factors for the development of CHD in children were smoking and alcoholism of parents, infectious and inflammatory diseases of the mother during pregnancy, as well as the associated use of antibacterial drugs; pregnancy-related factors as severe toxicosis, maternal age, gestosis of pregnant women, anemia; and socio-economic factors as marital status, low-income families, low educational level of the mothers of patients. The influence of these factors on the risk of CHD is indirect and largely depends on the quality of medical supervision of patients from socially disadvantaged families during pregnancy, since, as the study shows, such patients are registered late and follow medical recommendations less often.

Key words: Congenital heart defects, risk factors, demographic factors, biological factors, chemical factors pediatric cardiology

(Heart Vessels Transplant 2026; 10: 28-36. doi:10.24969/hvt.2026.629)

Introduction

Congenital heart diseases (CHDs) are structural abnormalities of the heart and/or great vessels that are present at birth (1-3). CHDs include an extensive range of defects that may be minor and require no intervention or may be life-limiting and require

complex surgery shortly after birth (4, 5). CHD is still the most common inborn defect with an approximate prevalence of 5–11 per 1000 live births and incidence of 1% (6).

CHD happen because the heart does not develop normally while the baby is growing in the womb.

Address for Correspondence: Sofia M. Shakhnabieva, Medical Faculty, Kyrgyz-Russian Slavic University, Bishkek, Kyrgyzstan

E-mail: shahnabieva_sofia@mail.ru

ORCID: Sofia M. Shakhnabieva - 0000-0002-6811-9580; Shakhshianam Sh. Khasanova- 0009-0008-0101-7505; Elena I. Kondrateva - 0000-0002-0674-4903.

Citation: Shakhnabieva SM, Khasanova ShSh, Sologubova TI, Kondrateva EI, Niiazalieva AD, Kurmanbakeev YM. Risk factors for the development of congenital heart defects in Kyrgyz Republic. Heart Vessels Transplant 2026; 10: 28-36. doi:10.24969/hvt.2026.629

Received: 30.01.2026 **Revised:** 01.03.2026 **Accepted:** 01.03.2026

Copyright ©2026 Heart, Vessels and Transplantation

Highlights

- The parental risk factors for CHD are:
- Habits - smoking and alcoholism of parents;
- Infectious and inflammatory diseases, associated use of antibacterial drugs of the mother during pregnancy;
- Pregnancy-related: severe toxicosis, maternal age, gestosis of pregnant women, anemia; Socio-economic: marital status, low-income of families, and low educational level

Graphical abstract



Heart, Vessels and Transplantation

Risk factors for the development of congenital heart defects in Kyrgyz Republic

Sofia M. Shakhnabieva, Shakhsianam Sh. Khasanova, Tatiana I. Sologubova, Elena I. Kondrateva, Ainagul D. Niiazalieva, Yuri M. Kurmanbakeev, Bishkek, Kyrgyzstan

The aim was to study the risk factors for congenital heart defects, to identify risk groups in whom CHD may occur.

Methods

A survey of parents of children with CHD (the main group, n=392) and parents of healthy children (the control group, n=100).

An analysis of the clinical, anamnestic, and medical-social characteristics of parents of children in both groups.

Results

The parental risk factors for CHD are (main group vs. controls):

Smoking and alcoholism of parents: maternal smoking (4.1 (0.2) vs. 1.0 (0.2), $p < 0.001$), maternal alcoholism (1.5 (0.1) vs. 0.0, $p < 0.001$), paternal smoking (57.1 (1.6) vs. 39.0 (5.9), $p < 0.01$) and paternal alcoholism (28.6 (0.3) vs. 2.0 (0.5) $p < 0.001$).

Infectious, inflammatory diseases and associated use of antibacterial drugs of the mother during pregnancy: intrauterine infections (20.1 (1.0) vs. 1.0 (0.2), $p < 0.001$) and ARVI (40.8 (1.5) vs. 12.0 (2.6); $p < 0.001$).

Pregnancy-related: severe toxicosis, maternal age, gestosis of pregnant women, anemia.

Socio-economic: marital status, low-income of families - (28.8 (1.3) vs 7.0 (1.6), $p < 0.001$) and low educational level of the mothers of patients (9.4 (0.5) vs. 7.0 (1.6), $p < 0.001$).

Conclusions

Thus, the most important risk factors for the development of CHD in children were smoking and alcoholism of parents, infectious and inflammatory diseases of the mother during pregnancy, as well as the associated use of antibacterial drugs; pregnancy-related factors and socio-economic factors as marital status, low-income families, low educational level of the mothers of patients. The influence of these factors on the risk of CHD is indirect and largely depends on the quality of medical supervision of patients from socially disadvantaged families during pregnancy, since, as the study shows, such patients are registered late and follow medical recommendations less often.

<http://hvt-journal.com/articles/art629>

doi:10.24969/hvt.2026.629

Genetics plays an important role in the formation of CHD (7, 8). It is common for CHD to occur because of changes, or mutations in the child's deoxyribonucleic acid (DNA) (2). The mutations in the DNA may or may not have come from the parents (9, 10).

Rarely, CHD are caused by particular genes that are inherited from the parents. That means a parent who has a CHD may be more likely to have a child with the defect (11-13). It is also known that fetuses with a prenatal diagnosis of CHD have an

increased risk of several comorbidities, including intrauterine growth restriction (14).

Previously, similar studies were conducted in Kyrgyzstan, but they did not include various groups of risk factors, and they were not comprehensive, so to undertake study on risk factors of CHD is timely and relevant.

The aim was to study the risk factors for congenital heart defects, to identify risk groups in whom CHD may occur.

Methods

Study design and population

This is a cross-sectional survey study. To study the main risk factors for having a child with CHD, parents of children with CHD (the main group, n=392) and parents of healthy children (the control group, n=100) were surveyed, followed by an analysis of the clinical, anamnestic, and medical-social characteristics of parents of children in both groups.

The survey was conducted in 19 family medicine centers from January 2024 to January 2025 in Bishkek, Kyrgyzstan among the main group of 392 patients with CHD: 120 patients with CHD (from 18 years old), 272 parents of children with CHD (under 18 years old), according to questionnaires developed by us in Russian and Kyrgyz languages. Inclusion criteria were: age 1 month - 60 years, diagnosed with «Congenital heart disease» (anatomical abnormalities of the heart/blood vessels), signed informed consent. Exclusion criteria: 1) severe concomitant pathology: diseases that are a contraindication to cardiac surgery; 2) absence of indications: defects that do not require active treatment or correction; 3) chromosomal/extracardial disorders: severe genetic syndromes that limit life if it is not a subject of study; 4) other reasons: infectious diseases or refusal to participate.

The data of 100 parents of healthy children without heart murmur who were evaluated in the children's department of maternity hospital No. 1 in Bishkek were used as a comparison group. The condition at birth of all children was assessed on the Apgar scale of at least 9 points, without signs of disability, having the first group of health.

Informed consent to participate in survey was signed by the parents/legal representatives and the patient (if >18 years old). The study was conducted according to the rules of the Helsinki 2024 agreement and approval of Institutional Ethics Committee was obtained.

Risk factors of CHD

We collected the following data:

- Demographic factors (gender, age, marital status, education level);
- Social and hygienic factors (material and living conditions in the family, bad habits (smoking, alcoholism, drug addiction, working at a personal computer, using a smartphone);
- Biological factors:
 - Diseases in close relatives (diseases of the cardiovascular system, a history of congenital anomalies, CHD in the immediate family);
 - Obstetric history (number of pregnancies, abortions, childbirths, features of the course of a real pregnancy (toxycosis, gestosis, bleeding, fetal infections));
 - Diseases, mothers suffered during pregnancy (acute respiratory viral infection (ARVI), anemia,

pyelonephritis, vaginitis, urethritis, colpitis, hypertension, diabetes, infection);

- Chemical factors (taking medications during pregnancy).

Survey

The survey was conducted through specially designed questionnaires in Russian and Kyrgyz languages. The questionnaires were distributed among the 19 Bishkek city family medicine centers, and they were handed over to pediatricians and internal medicine physicians, where patients live at the district and are monitored by these doctors. Parents of patients with CHD answered the questions. Then we collected the questionnaires and analyzed them. We did not interview patients from the country's regions.

During the survey, World Health Organization (WHO) data were used to determine smoking: about 200 puffs of smoke from smoldering tobacco leaves per day, which is 1 pack of cigarettes per day (15). Alcoholism was defined as a clinically established diagnosis, according to WHO: "a primary, chronic disease characterized by impaired control over alcohol intake, addiction to alcohol, alcohol consumption, despite the negative consequences." (16).

To determine family income, the average subsistence minimum in the Kyrgyz Republic was used (17), amounting to 8 758,6 soms (for 2025), low-income families had incomes in the family below the average subsistence minimum, middle-income families had incomes at the level of the average subsistence minimum, and families with high incomes were defined as having incomes above the average subsistence minimum.

When we studied the use of various medications, it turned out that medications were taken at different stages of pregnancy. Most of the women took various preventive medications before the 13th week of pregnancy (Iodomarin, folic acid, multivitamins).

Those who had iron deficiency anemia took iron supplements throughout pregnancy. The women took actovegin, calcium gluconate, antibiotics, nonsteroidal anti-inflammatory agents, and other medications in courses, depending on when the need arose.

Statistical analysis

Statistical data processing was performed using Microsoft Excel spreadsheets and the Statistica for Windows v. 10.0 application software package from StatSoft Inc. (USA), as well as the Biostat statistical program. Statistical processing of the materials was carried out using modern parametric and nonparametric methods of statistical analysis of biomedical research, as well as criteria of evidence-based medicine in accordance with modern requirements for the analysis of medical data. The representativeness was determined according to the generally accepted methodology of A.M. Merkov (1974) with an unknown number of the general population.

The statistical method of the study included the calculation of intensive, extensive indicators, the error of representativeness ($\pm m$), the criterion of reliability (t), the effectiveness of error-free prediction ($p=95.0\%$; $p=99.9\%$, with $p<0.01$; $p<0.001$), dynamic series (absolute increase, growth rate), correlation.

Results

The level of wealth and marital status in the families of children with CHD in the study groups is presented in Table 1.

Table 1. The level of wealth and marital status in families of children with CHD (per 100 respondents) in the study groups

#	Level of wealth and marital status of parents	Children with CHD Main group (n= 392)		Healthy children Control group (n= 100)		t/p
		the abs. number	P (m)	the abs. number	P (m)	
1.	High income	36	9.2 (0.5)	12	12.0 (2.7)	1.5/ >0.05
2.	Medium income	243	62.0 (1.5)	81	81.0 (1.0)	23.8/ <0.001
3.	Low income	113	28.8 (1.3)	7	7.0 (1.6)	20.5/ <0.001
4.	Married	221	56.4 (1.6)	87	87.0 (2.8)	11.8/ <0.001
5.	Not married	156	39.8 (1.5)	13	13.0 (2.8)	10.7/ <0.001
6.	No answer	15	3.8 (0.2)	-	-	-

P (m) is an intensive indicator and an error of representativeness; t is reliability; p is the effectiveness of an error-free forecast (with $p=95.0\%$, 99.9%) abs. – absolute, CHD - congenital heart disease

As follows from the data presented in Table 1, children in families with unfavorable socio-hygienic factors, such as family poverty ($p<0.001$), parents are not married ($p<0.001$), poor living conditions in the family were significantly more likely to suffer from CHD. At the same time, there was no increase in the incidence of CHD in families with good conditions ($p>0.05$).

The level of education and employment of parents in children with congenital heart disease is presented in Table 2.

When studying the influence of the level of education and employment of parents on the risk of developing CHD in

children, the following results were obtained. It turned out that the father's higher education ($p<0.001$) had a protective effect and reduced the risk of developing CHD.

A low level of education in the mother increased the risk of developing CHD in the child ($p<0.001$). The employment of the father reduced the risk of developing CHD in the child ($p<0.001$). The employment of the mother did not have a significant impact on the risk of developing CHD in children ($p>0.05$) (Table 2).

Table 2. The level of education and employment of parents in children with CHD

Variables	Categories of education and employment	Children with CHD Main group (n= 392)		Healthy children Control group (n= 100)		t/p
		the abs. number	P (m)	the abs. number	P (m)	
Father's education	Average	203	51.8(1.6)	33	33.0(5.5)	2.2/ <0.01
	Higher	126	32.1(1.4)	67	67.0(5.5)	4.3/ <0.001
	Primary	57	14.5(0.8)	0	-	-
Mother's education	Average	215	54.8(1.6)	58	58.0(6.0)	>0.05
	Higher	140	35.7(1.5)	35	35.0 (5.7)	>0.05
	Primary	37	9.4(0.5)	7	7.0(1.6)	3.4/ <0.001
Father's employment	Yes	298	76.0(1.1)	86	86.0(3.0)	4.0/ <0.001
	No	82	20.9(1.0)	14	14.0(3.0)	2.3/ <0.01
Mother's employment	Yes	137	34.9(1.4)	39	39.0(5.9)	>0.05
	No	255	65.0(1.5)	61	61.0(5.9)	>0.05

P (m) is an intensive indicator and an error of representativeness; t is reliability; p is the effectiveness of an error-free forecast (with $p=95.0\%$, 99.9%) abs. – absolute, CHD - congenital heart disease

Data on mother's age and cardiovascular disease in relatives of patients with CHD are presented in Table 3.

Table 3. Mother's age during pregnancy, cardiovascular diseases in close relatives of patients with CHD							
#	Variables	Children with CHD Main group (n= 392)		Healthy children Control group (n= 100)		Reliability	Interconnection (correlation)
		the abs. number	P (m)	the abs. number	P (m)	t/p	r _{xy}
Mother's age, years							
1.	From 16 to 20	45	11.5 (0.6)	13	13.0(2.8)	*/ >0.05	The connection is direct, strong +1,0
2.	21-30	182	46.4(1.6)	61	61.0(5.9)	*/ >0.05	
3.	From 31 to 45	165	42.1(1.5)	26	26.0(4.8)	6.3 /<0.001	
Cardiovascular diseases in close relatives of patients with CHD							
4.	No	272	69.4(1.3)	78	78.0(4.3)	*/>0.05	No connection
5.	Yes	120	30.6(1.3)	22	22.0(4.3)	*/ >0.05	

P (m) is an intensive indicator and an error of representativeness; t is reliability; p is the effectiveness of an error-free forecast (with p=95.0%, 99.9%)* – there is no significant difference, r_{xy} is a correlation relationship.
 abs. – absolute, CHD - congenital heart disease

From the data presented in Table 3, it follows that the age of the mother has a significant impact on the risk of CHD in children. In particular, the age of the mother over 30 years increases the likelihood of developing CHD in a child (p<0.001). The correlation is direct and strong, +1.0. At the same time, there

was no increase in the risk of having children with CHD in mothers less than 20 years of age (Table 3). It should also be noted that burdened inheritance of cardiovascular diseases from both the mother and father does not significantly increase the risk of having a child with CHD (p>0.05).

Table 4. Prevalence and influence of risk factors in parents on the development of CHD							
#	Variables	Children with CHD Main group (n= 392)		Healthy children Control group (n= 100)		p	Interconnection (correlation) r _{xy}
		Main group		Control group			
		the abs. number	P (m)	the abs. number	P (m)		
1.	Mother's smoking	16	4.1(0.2)	1	1.0(0.2)	<0.001	The connection is direct, strong (+1,0)
2.	Father's smoking	224	57.1(1.6)	39	39.0(5.9)	<0.01	
3.	Mother's alcoholism	6	1.5 (0.1)	0	0.0	<0.001	
4.	Father's alcoholism	112	28.6(0.3)	2	2.0(0.5)	<0.001	
5.	A mother's job on a PC	123	31.4(1.3)	11	11.0(2.4)	* >0.05	
6.	Phone usage (a cell phone (or mobile phone)	280	71.4 (1.3)	71	71.0(5.1)	* >0.05	

P (m) is an intensive indicator and an error of representativeness; t is reliability; p is the effectiveness of an error-free forecast (with p=95.0%, 99.9%)* – there is no significant difference, r_{xy} is a correlation relationship
 abs. – absolute, CHD - congenital heart disease, PC – personal computer

When assessing the effect of bad habits in parents on the risk of developing CHD in a child (Table 4), it turned out that maternal smoking (p<0.001), maternal alcoholism (p<0.001), paternal smoking (p<0.01), paternal alcoholism (p<0.001) significantly increases the risk of CHD in children.

The correlation is direct, strong, +1.0. At the same time, the mother's work on a personal computer (PC) and the use of a cell phone did not significantly affect the risk of developing CHD (p>0.05) (Table 4).

Table 5. Obstetric history and features of pregnancy in women of the main and control groups (per 100 respondents)

N	Obstetric history	Mothers with a risk factor (n= 392)				p	Interconnection (correlation) r_{xy}
		Main group		Control group			
		abs. number	P (m)	abs. number	P (m)		
1.	Abortions	112	28.6(1.3)	27	27.0(4.9)	* >0.05	The connection is direct, strong (+1,0)
2.	Severe toxicosis	275	70.2(1.4)	48	48.0(6.2)	<0.001	
3.	ARVI	160	40.8(1.5)	12	12.0(2.6)	<0.001	
4.	Gestosis	148	37.8(1.5)	13	13.0(2.3)	<0.001	
5.	IUI	79	20.1(1.0)	1	1.0(0.2)	<0.001	
6.	Bleeding	27	6.9(0.4)	6	6.0(1.4)	* >0.05	
7.	Anemia	49	12.5(0.7)	3	3.0(0.7)	<0.001	
8.	Pyelonephritis	47	12.0(0.7)	6	6.0(1.4)	<0.001	
9.	Vaginitis	3	0.8(0.05)	1	1.0(0.2)	<0.01	
10.	Urethritis	4	1.0(0.06)	1	1.0(0.2)	* >0.05	
11.	Colpitis	33	8.4(0.5)	4	4.0(1.0)	<0.001	

P (m) is an intensive indicator and an error of representativeness; t is reliability; p is the effectiveness of an error-free forecast (with $p=95.0\%, 99.9\%$)* – there is no significant difference, r_{xy} is a correlation relationship
 abs. – absolute, ARVI - acute respiratory viral infections, IUI - Intrauterine infection

An analysis of obstetric history data showed that the presence of previous abortions and the number of pregnancies had no significant effect on the risk of CHD in children ($p>0.05$) (Table 5). The most pronounced effect on the risk of developing CHD in a child was exerted by pathological conditions that develop during pregnancy. Thus, the presence of intrauterine infection (IUI) significantly increased the risk of CHD in children (

$p<0.001$), as well ARVI disease ($p<0.001$), anemia ($p<0.001$), gestosis ($p<0.001$), and severe toxicosis ($p<0.001$). The correlation is direct and strong (+1.0). At the same time, we did not find an association between the presence of inflammatory diseases of the urinary tract and bleeding in the mother with an increased risk of CHD in children ($p>0.05$) (Table 5).

Table 6. Frequency of taking medications during pregnancy (per 100 respondents)

N	Drugs	Frequency of taking the drug				t/p
		Main group		Control group		
		abs. number	P (m)	abs. number	P (m)	
1.	Iron-containing preparations	35	8.9(0.5)	13	13.0(2.8)	2.0 /<0.01
2.	Iodomarin	12	3.1(0.2)	3	3.0(0.7)	*/>0.05
3.	Multivitamins	23	5.9(0.3)	4	4.0(1.0)	7.0/<0.001
4.	Actovegin	3	0.8(0.05)	2	2.0(0.5)	20.0/ <0.001
5.	Calcium gluconate	8	2.0(0.1)	1	1.0(0.2)	10.0/ <0.001
6.	Folic acid	9	2.3(0.1)	3	3.0(0.7)	10.0/ <0.001

P (m) is an intensive indicator and an error of representativeness; t is reliability; p is the effectiveness of an error-free forecast (with $p=95.0\%, 99.9\%$)* – there is no significant difference, r_{xy} is a correlation relationship

A study was conducted on the effect of pregnant women taking a number of medications (iron-containing, iodomarin, multivitamins, folic acid, calcium gluconate, and actovegin) on the risk of CHD in children. It turned out that taking multivitamins ($p<0.001$) and calcium gluconate ($p<0.001$) increases the risk of CHD. Taking iron-containing drugs, folic acid, and actovegin was not associated with an increased

risk of having children with developmental abnormalities of the cardiovascular system. However, these data should be interpreted with caution due to the small number of such observations in both the main and control groups (Table 6).

At the same time, a relationship has been established between the birth of children with CHD and taking medications for infections during pregnancy (Table 7).

Table 7. Frequency of medications taken for acute respiratory viral infections during pregnancy (per 100 respondents)

N	Medications	Frequency of taking the drug				t/p
		Main group		Control group		
		abs. number	P (m)	abs. number	P (m)	
1.	Antipyretics	30	7.7(0.4)	8	8.0(1.8)	*/>0.05
2.	Expectorants	19	4.8(0.3)	2	2.0(0.5)	10.0/<0.001
3.	Antihistamines	8	2.0(0.1)	-	-	-
4.	Antiviral drugs	13	3.3(0.2)	-	-	-
5.	NSAIDs	23	5.9(0.3)	1	1.0(0.2)	16.3/<0.001
6.	Antifungal	14	3.6(0.2)	-	-	-
7.	Antibacterial	107	27.3(1.3)	4	4.0(1.0)	25.8/<0.001

P (m) is the intensive indicator and the error of representativeness; T is the reliability; P is the effectiveness of an error-free forecast (at p=95,0%, 99,9%), * – there is no significant difference
 NSAIDs – nonsteroidal anti-inflammatory drugs

Thus, the use of antibacterial agents by women for intrauterine infections (p<0.001), nonsteroidal anti-inflammatory drugs (p<0.001), expectorants was associated with an increased risk of childbirth with the CHD. Antibacterial drugs taken by pregnant women were of various groups, such as penicillins (ampicillin, amoxicillin, amoxiclav), cephalosporins (cefazolin, cefuroxime), macrolides (erythromycin, roxithromycin, clarithromycin, azithromycin), tetracyclines (doxycycline), fluoroquinolones (ciprofloxacin, norfloxacin). There was no increase in the risk of CHD in children when taking antipyretic, antiviral, antifungal, and antihistamines (p>0.05) (Table 7).

Therefore, the main risk factors for the development of CHD in children were the bad habits of parents (the correlation is direct, strong, +1.0); infectious and inflammatory diseases of the mother during pregnancy, such as ARVI and intrauterine infections, as well as the associated use of antibacterial drugs. A significant contribution to the risk determination was made by conditions directly related to pregnancy (severe toxicosis, maternal age, gestosis of pregnant women, anemia (the correlation is direct, strong, +1.0)). Attention was drawn to the association of the incidence of CHD with a number of socio-economic factors: marital status, low-income families, low educational level of parents, primarily the mother.

Discussion

Thus, according to the data of the screening questionnaire cards of the main group, consisting of parents of patients with CHD and the comparison group, which included parents of healthy children, the role of medical and social factors on the part of the father and mother was determined.:

1. Demographic factors as the age of the mother over 30 years increases the likelihood of developing CHD in a child (p<0.001). The correlation is direct, strong, +1.0. When parents are not married (p<0.001) and the low level of education of the mother (p<0.001) increased the risk of CHD in the child).

2. Social and hygienic factors: children in low-income families were significantly more likely to suffer from CHD (p<0.001). Maternal smoking (p<0.001), maternal alcoholism (p<0.001), paternal smoking (p<0.01), paternal alcoholism (p<0.001) increase the risk of having a child with CHD. The correlation is direct, strong, +1.0).
3. Biological factors like diseases: presence of IUI, ARVI disease, anemia, gestosis, severe toxicosis significantly increased the risk of CHD in children (correlation is direct, strong, +1.0).
4. Chemical factors: taking multivitamins and calcium gluconate increases the risk of CHD. The use of antibacterial agents by women for IUI, nonsteroidal anti-inflammatory drugs, expectorants was associated with an increased risk of having children with CHD.

Our survey results are confirmed by scientific research in other countries. Population-based studies in Ontario, Canada, have shown that mothers living in deprived neighborhoods had 27-29% higher chances of having infants with CHD compared to those in wealthier areas (18). The etiology of CHD is known to be multifactorial, involving both genetic predisposition and environmental influences. Low socioeconomic status is recognized as a risk factor (19).

Lower maternal income, poverty, lower educational level, and unemployment status all showed positive associations with CHD risk. Mothers in the lowest income neighborhoods had 15-29% higher odds of having children with CHD compared to those in the highest income neighborhoods (20).

If we compare our results with data from other researchers in the world, it turned out that nutritional factors, and prenatal care access – may play important roles. Maternal comorbidities explain only about 13% of the relationship between social deprivation and CHD incidence, suggesting that other pathways – potentially including environmental

exposures, nutritional factors, and prenatal care access – may play important roles (21).

The revealed data on the pathological course of the present pregnancy, as risk factors for the development of fetal CHD, should be used during the registration of a woman in a women's clinic.

Taking into account all these factors make it possible to determine and prevent the risk of having a child with CHD at an early stage, and will allow for the adjustment of the management plan for women in the periconceptual period.

Study limitations

The study's main limitation is that it was conducted only in Bishkek, and did not include other cities and rural areas. Further studies on risk factors of CHD in children residing rural areas should be thought.

Another limitation is that it is a survey and cross-sectional study. To confirm associations, the prospective studies should be done.

Conclusion

Thus, the most important risk factors for the development of CHD in children were bad habits of parents, primarily smoking and alcoholism, infectious and inflammatory diseases of the mother during pregnancy, acute respiratory viral infections and intrauterine infections, as well as the associated use of antibacterial drugs. Conditions directly related to pregnancy (severe toxicosis, maternal age, gestosis of pregnant women, anemia) made a less noticeable but significant contribution to determining the risk. The increased risk of having a child with CHD was determined by a number of socio-economic factors: marital status, low-income families, low educational level of the mothers of patients. The influence of these factors on the risk of CHD is indirect and largely depends on the quality of medical supervision of patients from socially disadvantaged families during pregnancy, since, as the study shows, such patients are registered late and follow medical recommendations less often.

Ethics: Informed consent to participate in survey was signed by the parents/legal representatives and the patient (if >18 years old). The study was conducted according to the requirements of the Helsinki 2024 declaration and approval of Institutional Ethics Committee was obtained.

Peer-review: External and internal

Conflict of interest: None to declare

Authorship: S.M.Sh., Sh.Sh.Kh., T.I.S., E.I.K., A.D.N., and Y.M.K. equally contributed to preparation of manuscript and fulfilled all authorship criteria

Acknowledgements and funding: None to declare

Statement on A.I.-assisted technologies use: The authors did not use AI-supported technologies in the preparation of this manuscript.

Data and material availability: Does not apply

References

1. Wilsdon A, Loughna S. Human genetics of congenital heart defects. *Adv Exp Med Biol* 2024; 1441: 57-75.
2. Meisner JK, Martin DM. Congenital heart defects in CHARGE: The molecular role of CHD7 and effects on cardiac phenotype and clinical outcomes. *Am J Med Genet* 2020; 184: 81-9. doi: 10.1002/ajmg.c.31761
3. Guijarro C, Kelly RG. On the involvement of the second heart field in congenital heart defects. *C R Biol* 2024; 347: 9-18. doi: 10.5802/crbio.151
4. Palma A, Morais S, Silva PV, Pires A. Congenital heart defects and preterm birth: Outcomes from a referral center. *Rev Port Cardiol* 2023; 42: 403-10. doi: 10.1016/j.repc.2022.05.009
5. Diaz Kane MM. Pulse oximetry screening for congenital heart defects in the newborn nursery: A review for the general pediatrician. *Pediatr Ann* 2022; 51: e411-3. doi: 10.3928/19382359-20220913-01
6. Micheletti A. Congenital heart disease classification, epidemiology, diagnosis, treatment, and outcome. In: Flocco, S, Lillo A, Dellafiore, F, Goossens E. (eds) *Congenital Heart Disease*. Springer: Cham, Switzerland 2019; 303p. https://doi.org/10.1007/978-3-319-78423-6_1
7. Houyel L, Meilhac SM. Heart development and congenital structural heart defects. *Annu Rev Genomics Hum Genet* 2021; 22: 257-84. doi: 10.1146/annurev-genom-083118-015012
8. Arvind B, Saxena A. Timing of interventions in infants and children with congenital heart defects. *Indian J Pediatr* 2020; 87: 289-94. doi: 10.1007/s12098-019-03133-w
9. Kathiriya IS. Insights into human heart development and congenital heart defects. *Development* 2024; 151: dev204302. doi: 10.1242/dev.204302
10. Perlow G, Kochilas L. Long-term outcomes after interventions for infants with congenital heart defects. *Clin Perinatol* 2025; 52: 761-70. doi: 10.1016/j.clp.2025.08.007
11. Barbosa M, Jørgensen FS, Christiansen M, Steensberg JN, Ekelund CK, Vedel C. Prenatal screening for severe congenital heart defects in Denmark. *Ugeskr Laeger* 2025; 187: V11240763. doi: 10.61409/V11240763
12. Bourdon G, Lenne X, Godart F, Storme L, Theis D, Subtil D, et al. Epidemiology of congenital heart defects in France from 2013 to 2022 using the PMSI-MCO (French Medical Information System Program in Medicine, Surgery, and Obstetrics) database. *PLoS One* 2024; 19: e0298234. doi: 10.1371/journal.pone.0298234
13. Miao H, Zeng Q, Shi Z, Xia Y, Shi L, Chen D, et al. Spatio-temporal distribution and demographic characteristics of congenital heart defects in Guangdong, China, 2016-2020. *Front Public Health* 2022; 10: 813916. doi: 10.3389/fpubh.2022.813916

14. Ferreira BD, Barros T, Moleiro ML, Guedes-Martins L. Preeclampsia and fetal congenital heart defects. *Curr Cardiol Rev* 2022; 18: 80-91. doi: 10.2174/1573403X18666220415150943
15. World Health Organization. Tobacco. Available at: URL: <https://who.int>
16. World Health Organization. Alcohol use. Available at: URL: <https://who.int>
17. National Statistical Committee of the Kyrgyz Republic. Living standard. Available at: URL: <https://stat.gov.kg>
18. Miao Q, Dunn S, Wen SW, Lougheed J, Yang P, Davies M, et al. Association between maternal marginalization and infants born with congenital heart disease in Ontario Canada. *BMC Public Health* 2023; 23: 790.
19. Diller GP, Arvanitaki A, Opotowsky AR, Jenkins K, Moons P, Kempny A, et al. Lifespan perspective on congenital heart disease research: JACC State-of-the-Art Review. *J Am Coll Cardiol* 2021; 77: 2219-35.
20. Miao Q, Dunn S, Wen SW, Lougheed J, Maxwell C, Reszel J, et al. Association of maternal socioeconomic status and race with risk of congenital heart disease: a population-based retrospective cohort study in Ontario, Canada. *BMJ Open* 2022; 12: e051020.
21. Peyvandi S, Bayer RJ, Chambers CD, Norton ME, Rajagopal S, Ryckman KK, et al. Environmental and socioeconomic factors influence the live-born incidence of congenital heart disease: A population-based study in California. *J Am Heart Assoc* 2020; 9: e015255.